



GIG
CYMRU
NHS
WALES

Shingles vaccination for those aged 70 to 79



Shingles affects a lot of people as they get older.

It often causes severe pain which can last for months or sometimes years.

Shingles is more common and more severe the older you are.

There is a vaccine to help protect older people from shingles and the long term pain that can follow.

OGL

© Crown Copyright 2015
Print ISBN 978 1 4734 3442 4
Digital ISBN 978 1 4734 3443 7
WG25207

What is shingles?

Shingles (also known as herpes zoster) is an infection of a nerve and the skin around it. It usually causes a painful rash of fluid-filled blisters. The blisters burst and turn into sores. The rash usually affects an area on just one side of the body, most commonly the chest but sometimes the head, face and eye.

How long does it last and how serious can it be?

The rash usually appears a few days after the pain starts, and the blisters crust over in about a week but can last longer.

Most people recover fully, but for some the pain can go on for several months and sometimes lasts for years. The older you are the more likely you are to have long-lasting pain. This is called post-herpetic neuralgia (PHN) and can cause a severe burning, throbbing or stabbing pain. Current treatments for PHN are not very effective.

Sometimes shingles affects the eye or the skin around the eye. This can cause severe pain or even blindness.

What causes shingles?

Shingles is caused by the same virus as chickenpox – varicella zoster virus.

When you recover from chickenpox, most of the virus is destroyed but some survives and lies inactive in the body. This virus can reactivate later in life when your immune system is weakened by things such as age, stress, mental health problems or drugs that reduce immune response.

How do you catch shingles?

You don't catch shingles. Chickenpox virus caught earlier in life reactivates later to cause shingles. However, the fluid in shingles blisters can cause chickenpox by infecting someone who has never had it (usually a child).

How common is shingles?

Almost everyone gets chickenpox and about one in five people who have had it go on to get shingles.

Every year in England and Wales, tens of thousands of people will get shingles. It is most common in people aged 70 years and over, and of these:

- about 14,000 go on to develop PHN and
- over 1,400 are admitted to hospital.

Around one in 1,000 people over 70 who gets shingles dies of the infection.

How effective is the vaccination?

Shingles vaccination will reduce your chances of developing shingles by more than a third. If you do go on to have shingles it will probably be a milder and shorter illness.

Where is the vaccination given and will I need one every year?

Most people will have the vaccine at their GP surgery.

The vaccine will be given in your upper arm. You only need one vaccination, you do not need it every year.

Will there be any side effects?

Side effects are usually quite mild and don't last very long. The most common side effects are headache and redness, pain, swelling, itching, warmth or bruising at the site of the injection. If the side effects last for more than a few days you should discuss this with your GP or practice nurse.

Should I take any precautions after vaccination?

Rarely a rash of small blisters develops where the injection was given. The rash should be covered, and until it crusts over, you should avoid contact with anyone with

a weakened immune system, newborn babies or who is pregnant and has never had chickenpox.

How safe is the vaccine – has it been used in other countries?

The shingles vaccine has been thoroughly tested and meets UK and European safety and licensing requirements. It has been used across the UK for those aged 70 and 79 from 2013, and used in other countries including the United States of America and Canada.

Who will get the vaccine?

The routine shingles vaccination programme was introduced for all 70 year olds in 2013. As people become 70 they become eligible for the vaccine and can have it at any time before their 80th birthday.

You can have the vaccine if you:

were born on or after 02 September 1942.

AND

have already reached the age of 70 years.

AND

are under 80 years of age.

There is also a 'catch up' campaign.

You can have the vaccine if you are aged 78 or 79 years old.

What about people outside the 70 to 79 age group?

Shingles is less common in younger people and those aged under 70 will not routinely get the vaccine.

The vaccine is less effective as people get older and people aged 80 and over will not be routinely vaccinated.

Each year more people will become eligible to have a shingles vaccine until everyone between the ages of 70 and 79 can have it.

Do I need to do anything to get the vaccination?

Your GP will invite you for the vaccination. You can have it at any time of year, including at the same time as your flu jab in the autumn. If you think you may have missed your invitation, contact your GP surgery.

Are there people who shouldn't have the vaccination?

People who have a weakened immune system, for example due to current cancer treatment, should not have the vaccine. Also anyone who has had a severe reaction to anything used in the vaccine shouldn't have it. Your GP or practice nurse will check this.

Where can I get more information?

You can find out more about the vaccine, including vaccine contents and potential side effects, at www.medicines.org.uk/emc/ by entering the name of the vaccine



in the search box and looking at the patient information leaflet (PIL).

The routine immunisation schedule that shows you which immunisations are offered in Wales is available from www.publichealthwales.org/routineimmunisation-schedule-wales

If you have any questions or want more information, talk to your doctor or nurse or call NHS Direct Wales on 0845 4647 or visit

www.nhsdirect.wales.nhs.uk/



To order more copies of this leaflet, email: hplibrary@wales.nhs.uk or telephone 08456 064050.

This leaflet is produced by:

Public Health Division
Welsh Government and the Vaccine
Preventable Disease Programme
Public Health Wales